

References:

1. M. A. Mansour, S. A. Bekheet, S. S. Al-Rejaie et al., "Ginger ingredients inhibit the development of diethylnitrosoamine induced premalignant phenotype in rat chemical hepatocarcinogenesis model," *BioFactors*, vol. 36, no. 6, pp. 483–490, 2010. [View at Publisher](#) · [View at Google Scholar](#) · [View at Scopus](#).
2. M. Wuthi-udomlert, W. Grisanapan, O. Luanratana, and W. Caichompoo, "Antifungal activity of *Curcuma longa* grown in Thailand," *Southeast Asian Journal of Tropical Medicine and Public Health*, vol. 31, no. 1, pp. 178–182, 2000.
3. D. Greenwood, *Microbiology, A Guide To Microbiology Infections: Pathogenesis, Immunity, Laboratory Diagnosis and Control*, Churchill Livingstone, Philadelphia, Pa, USA, 17th edition, 2007.
4. <http://www.turmericforhealth.com/turmeric-benefits/benefits-of-turmeric-essential-oil#ixzz3mzoQ1iYb>
5. Metabolism of ginger component [6]-shogaol in liver microsomes from mouse, rat, dog, monkey, and human. *Molecular Nutrition and Food Research*. January 2013.
6. Characterization of thiol-conjugated metabolites of ginger components shogaols in mouse and human urine and modulation of the glutathione levels in cancer cells by [6]-shogaol. *Molecular Nutrition and Food Research*, January 2013