

References :

1. Erickson, D.R., Pryde, E. H., Brekke, O.L., Mounts, T.L., and Falb, R. A. (Editors) 1980. Handbook of Soy Oil Processing and Utilization. American Soybean Association, St. Louis, MO and American Oil Chemists' Society, Champaign, IL.
2. Hendricks, J. D., Sinnhuber, R. O., Loveland, P. M., Pawlowski, N.E., and Nixon, J. E. 1980. Hepatocarcinogenicity of glandless cottonseeds and cottonseed oil to rainbow trout (*Salmo gairdnerii*). Science 208, 309-311.
3. Harris, J. A., Magne, F. C., and Skau, E. L. 1964. Methods for the determination of cyclopropenoid fatty acids. IV. Application of the step-wise HBr titration method to the analysis of refined and crude cottonseed oils. J. Am. Oil Chem. Soc. 41, 309-311.
4. Reddy, N. R., Sathe, S.K., and Salunkhe, D.K. 1982C. Phytates in legumes and cereals. Adv. Food Res. 28, 1-92.
5. Berardi, L. C., and Goldblatt, L. A. 1980. Gossypol. In Toxic Constituents of Plant Foodstuffs. 2nd Edition, pp. 183-237. I.E. Liener (Editor). Academic Press, New York.
6. Lindsey, T.O., Hawkins, G.E., and Guthrie, L.D.1980. Physiological responses of lactating cows to gossypol from cottonseed meal rations. J. Dairy Sci. 63, 562-573.
7. Rosen, G. D. 1958. Groundnuts (peanuts) and groundnut meal. In Processed Plant Protein Foodstuffs, pp. 419-468. A. M. Altschul (Editor). Academic Press, New York.